CO1.2: Life expectancy at birth

Definitions and methodology

This indicator uses two measures to capture life expectancy:

- Life expectancy at birth, defined as the average number of years a new-born
 child would live if prevailing patterns of mortality at the time of its birth were to
 stay the same throughout their life.
- Health-adjusted life expectancy (HALE) at birth, defined as the average number of years that a new-born child can expect to live in "full health" after taking into account years expected to be lived in less than full health due to disease and/or injury.


Data on life expectancy at birth comes either from OECD Health Statistics or from the World Bank, while data on HALE at birth comes from the World Health Organization (WHO) Global Health Observatory, which updates its data every 5 years.

Key findings

Current life expectancies for new-born children vary considerably across the covered Asia/Pacific countries (Chart CO1.2.A). In five of the covered countries (Australia, Japan, Korea, Singapore, and New Zealand), current life expectancies are relatively high. In these countries, new-born children can expect to live at least until the age of 82 – above the average for OECD countries (80). However, current life expectancies at birth are lower in China, Malaysia, Thailand, and Viet Nam – in these countries, new-born children can expect to live until around age of 75-78 – and are much lower in Mongolia and Indonesia at about 70 years.

Chart CO1.2.A. Life expectancy at birth by gender, 2021

Average number of years a new-born infant would live if prevailing patterns of mortality at the time of its birth were to stay the same throughout its life



a) The OECD average refers to the unweighted average across OECD member countries with available and comparable data in 2021. See OECD Family Database Indicator CO1.2 (http://www.oecd.org/els/family/database.htm) for more detail.

Sources: Australia, Japan, New Zealand, Indonesia, and China: OECD Health Statistics; OECD average: OECD Family Database Indicator CO1.2; Malaysia, Thailand and Viet Nam: World Bank (2023), World Development Indicators; Mongolia: Statistical Yearbook 1991-2021; Singapore: Department of Statistics Singapore (2022). Life Expectancy by Sex.

Other relevant indicators: SF2.1 Fertility rates; CO1.1 Infant mortality; CO1.3 Low birth weight; CO1.4 Vaccination rates.

1 Updated: December 2023

Family Database in the Asia-Pacific Region, http://oe.cd/fdb-asia OECD and OECD KOREA Policy Centre

Across countries, girls tend to have higher life expectancies than boys, though the extent of the gap varies (Chart CO1.2.A). The largest current gender gaps in life expectancy are in Mongolia, Thailand, and Viet Nam, where life expectancy at birth is about 7-9 years higher for girls than for boys: in Mongolia, a new-born girl can expect to live almost 9.5 years longer than a new-born boy. The smallest gender gap is in New Zealand at 3.5 years.

All Asia/Pacific countries included here seen considerable gains in life expectancies at birth over the past four or five decades (Table CO1.2.A). The current life expectancies at birth are at least 10 years longer than they were in 1960 (or in the earliest year with available data) for all covered countries, with the largest gains made in Korea (an increase of more than 20 years since 1970), Indonesia and Thailand (22 and 23 years since 1960), and especially China (45 years since 1960).

COVID-19 has had a major impact on life expectancy due to the exceptionally high number of deaths caused by the pandemic. Prior to the pandemic, life expectancy increased in all countries; however, life expectancy has stabilized since 2019 in many countries or declined (e.g. Indonesia and Viet Nam (See Table CO1.2.A and OECD 2023, Health at a Glance).

Table CO1.2.A. **Trends in life expectancy at birth, 1960-2022**Average number of years a new-born infant would live if prevailing patterns of mortality at the time of its birth were to stay the same throughout its life

| | 1960 | 1970 | 1980 | 1990 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 |
|----------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| OECD-31 av erage (a) | 68.3 | 70.0 | 72.5 | 74.8 | 77.1 | 78.4 | 79.7 | 80.6 | 80.9 | 80.9 | 81.0 | 81.3 | 80.8 | 80.7 | |
| Australia | 70.9 | 70.8 | 74.6 | 76.9 | 79.2 | 80.8 | 81.7 | 82.4 | 82.4 | 82.5 | 82.7 | 82.9 | 83.2 | 83.3 | |
| China | 33.3 | 59.1 | 63.9 | 67.8 | 71.4 | 73.7 | 75.3 | 76.7 | 77.0 | 77.2 | 77.2 | 77.7 | 78.0 | 78.1 | 78.2 |
| Indonesia | 46.5 | 52.6 | 58.2 | 62.7 | 66.2 | 65.8 | 68.5 | 69.5 | 69.7 | 69.8 | 69.9 | 70.3 | 70.5 | 68.8 | 67.6 |
| Japan | 67.8 | 72.0 | 76.1 | 78.9 | 81.2 | 82.0 | 82.9 | 83.9 | 84.1 | 84.2 | 84.3 | 84.4 | 84.7 | 84.5 | |
| Korea | | 62.3 | 66.2 | 71.7 | 76.0 | 78.2 | 80.2 | 82.1 | 82.4 | 82.7 | 82.7 | 83.3 | 83.5 | 83.6 | |
| Malaysia | 56.5 | 63.1 | 68.2 | 71.3 | 72.8 | 74.1 | 74.4 | 75.1 | 75.3 | 75.5 | 75.6 | 75.8 | 75.9 | 74.9 | |
| Mongolia | | | | | 63.2 | 65.2 | 68.1 | 69.9 | 69.6 | 69.9 | 70.2 | 70.4 | 70.7 | 71.0 | |
| New Zealand | | 71.5 | 73.2 | 75.5 | 78.4 | 79.8 | 80.8 | 81.7 | 81.7 | 81.9 | 81.8 | 82.1 | 82.3 | 82.3 | |
| Singapore | 65.7 | 68.3 | 72.2 | 75.3 | 78.0 | 80.1 | 81.7 | 82.9 | 83.0 | 83.2 | 83.4 | 83.7 | 83.7 | 83.5 | |
| Thailand | 54.7 | 59.4 | 64.4 | 70.3 | 70.6 | 72.1 | 73.9 | 75.1 | 75.3 | 76.7 | 76.9 | 77.2 | 77.3 | | |
| Viet Nam | 59.7 | 56.2 | 66.2 | 69.2 | 72.5 | 73.3 | 73.5 | 73.9 | 73.9 | 74.0 | 74.0 | 74.1 | 75.4 | 73.6 | |

a) The OECD-31 average refers to the unweighted average across the 38 OECD member countries, except for Canada, Israel, Korea, Latvia, Lithuania, Mexico, and Slovenia, with available and comparable data for the whole period. See OECD Family Database Indicator CO1.2 (http://www.oecd.org/els/family/database.htm) for more detail.

Sources: Australia, Japan, New Zealand, Indonesia, and China: OECD Health Statistics; OECD average: OECD Family Database Indicator CO1.2; Malaysia, Thailand and Viet Nam: World Bank (2023), World Development Indicators; Mongolia: Statistical Yearbook 1991-2021; Singapore: Department of Statistics Singapore (2022). Life Expectancy by Sex.


Life expectancy does not however provide a complete picture of the health status of the population, especially if extra years of life are not lived in good health. In Asia/Pacific countries as also in OECD countries, 'health-adjusted' life expectancy at birth – that is, the number of years a new-born infant can expect to leave in full health – is often much shorter than actual life expectancy, especially for new-born girls (Chart CO1.2.B). The smallest differences are in China, where 'health-adjusted' life expectancies at birth are 7.6 years shorter than actual life expectancies for boys and 9.2 years shorter for girls. The largest differences are in Australia – where the health-adjusted life expectancies at birth is 10.8 years shorter than the actual life expectancy for boys, and 13.3 years shorter for girls.

2 Updated: December 2023

Family Database in the Asia-Pacific Region, http://oe.cd/fdb-asia OECD and OECD KOREA Policy Centre

Chart CO1.2.B. Life expectancy at birth and Health-Adjusted Life Expectancy (HALE) at birth, 2019

Average number of years a new-born infant can expect to live if prevailing patterns of mortality at the time of its birth were to stay the same throughout its life (life expectancy at birth), and average number of years that a new-born infant can expect to live in "full health" by taking into account years lived in less than full health due to disease and/or injury (HALE)



a) The OECD average of Life expectancy refers to the unweighted average across OECD member countries with available and comparable data. See OECD Family Database Indicator CO1.2 (http://www.oecd.org/els/family/database.htm) for more detail.

Sources: WHO Global Health Observatory, 2019

Comparability and data issues

The data on life expectancy at birth shown in Chart CO1.2.A and Table CO1.2.A are taken from three sources: OECD Health Statistics for OECD member countries, the World Bank Open Data Database for Malaysia, Thailand and Viet Nam, and nationally published sources for all other countries. In all cases the data was originally collected from national statistical offices or from data published by other international organisations, such as the United Nations Population Division. It is possible that some of the international variation in life expectancy at birth may be due to differences between countries in the registering of deaths or recording of mortality rates (see here for more details on the data collected and published by OECD Heath Statistics). For the data published by OECD Health Statistics, life expectancy at birth for the total population is estimated by the OECD Secretariat using the unweighted average of life expectancies for men and women.

The data on life expectancy and Healthy Life Expectancy (HALE) at birth shown in Chart CO1.2.B are taken from the WHO Global Health Observatory, which itself calculates estimates based on life tables constructed by WHO using Sullivan's method. For more information on the data and methods used in the calculation of the HALE data, see the Global Health Observatory website.

Sources and further reading: OECD Health Statistics, http://www.oecd.org/els/health-systems/health-data.htm, World Health Organization Global Health Observatory, https://www.who.int/gho/en/; World Bank Open Data Database, https://data.worldbank.org/; OECD/WHO (2022), Health at a Glance: Asia/Pacific 2022: Measuring Progress Towards Universal Health Coverage, OECD Publishing, Paris, https://doi.org/10.1787/c7467f62-en.

3 Updated: December 2023